

# Coefficients of Chiral Perturbation Theory

Bing An Li

Department of Physics and Astronomy, University of Kentucky

Lexington, KY 40506, USA

## **Abstract**

Based on an effective chiral theory of pseudoscalar, vector, and axial-vector mesons, the coefficients of the chiral perturbation theory are predicted. There is no new parameter in these predictions.

The chiral perturbation theory(ChPT) proposed a decade ago[1] is successful in parametrizing low-energy  $QCD$ . The chiral symmetry revealed from  $QCD$ , quark mass expansion, and momentum expansion are used to construct the Lagrangian of ChPT

$$\begin{aligned}
\mathcal{L} = & \frac{f_\pi^2}{16} Tr D_\mu U D^\mu U^\dagger + \frac{f_\pi^2}{16} Tr \chi (U + U^\dagger) + L_1 [Tr (D_\mu U D^\mu U^\dagger)]^2 + L_2 (Tr D_\mu U D_\nu U^\dagger)^2 \\
& + L_3 Tr (D_\mu U D^\mu U^\dagger)^2 + L_4 Tr (D_\mu U D^\mu U^\dagger) Tr \chi (U + U^\dagger) + L_5 Tr D_\mu U D^\mu U^\dagger (\chi U^\dagger + U \chi) \\
& + L_6 [Tr \chi (U + U^\dagger)]^2 + L_7 [Tr \chi (U - U^\dagger)]^2 + L_8 Tr (\chi U \chi U + \chi U^\dagger \chi U^\dagger) \\
& - i L_9 Tr (F_{\mu\nu}^L D^\mu U D^\nu U^\dagger + F_{\mu\nu}^R D^\mu U^\dagger D^\nu U) + L_{10} Tr (F_{\mu\nu}^L U F^{\mu\nu R} U^\dagger). \tag{1}
\end{aligned}$$

The parameters in the chiral perturbation theory(1) are pion decay constant  $f_\pi$  and the 10 coefficients. These parameters are determined by fitting experimental data[1,2,3,4,5,6,7].

The chiral perturbation theory is rigorous and phenomenologically successful in describing the physics of the pseudoscalar mesons at low energies. Models attempt to deal with the two main frustrations that the ChPT is limited to pseudoscalar mesons at low energy and contains many coupling constants which must be measured. However, the chiral perturbation theory sets a low energy limit for all models. The price paid for including more mesons and determining the coefficients is to make additional assumption. Many models try to predict the chiral couplings(see Table 2). In Ref.[8] an extended Nambu-Jona-Lasinio model(ENJL)

$$\mathcal{L} = \bar{q} \{ \gamma^\mu (v_\mu + \gamma_5 a_\mu) - (s - ip\gamma_5) \} q$$

has been used to predict the coefficients of the ChPT. Nine of the 10 coefficients,  $L_{1,2,3,4,5,6,8,9,10}$ , are predicted in Ref.[8b] and listed in Table II.

I have proposed an effective chiral theory of pseudoscalar, vector, and axial-vector mesons[9,10]. It provides a unified description of low-lying meson physics. This effective theory is phenomenologically successful. It is natural to study whether the Lagrangian(1) of the chiral perturbation theory can be derived from this effective theory and the 10 coefficients can be predicted. The Lagrangian of this theory is

$$\begin{aligned} \mathcal{L} = & \bar{\psi}(x)(i\gamma \cdot \partial + \gamma \cdot v + \gamma \cdot a\gamma_5 - mu(x))\psi(x) - \bar{\psi}(x)M\psi(x) \\ & + \frac{1}{2}m_0^2(\rho_i^\mu \rho_{\mu i} + K^{*\mu} K_\mu^* + \omega^\mu \omega_\mu + \phi^\mu \phi_\mu + a_i^\mu a_{\mu i} + K_1^\mu K_{1\mu} + f^\mu f_\mu + f_{1s}^\mu f_{1s\mu}) \end{aligned} \quad (2)$$

where M is the quark mass matrix

$$\begin{pmatrix} m_u & 0 & 0 \\ 0 & m_d & 0 \\ 0 & 0 & m_s \end{pmatrix},$$

$v_\mu = \tau_i \rho_\mu^i + \lambda_a K_\mu^{*a} + (\frac{2}{3} + \frac{1}{\sqrt{3}}\lambda_8)\omega_\mu + (\frac{1}{3} - \frac{1}{\sqrt{3}}\lambda_8)\phi_\mu$ ,  $a_\mu = \tau_i a_\mu^i + \lambda_a K_{1\mu}^a + (\frac{2}{3} + \frac{1}{\sqrt{3}}\lambda_8)f_\mu + (\frac{1}{3} - \frac{1}{\sqrt{3}}\lambda_8)f_{1s\mu}$ , and  $u = \exp\{i\gamma_5(\tau_i \pi_i + \lambda_a K_a + \lambda_8 \eta_8 + \eta_0)\}$ .  $u$  can be written as

$$u = \frac{1}{2}(1 + \gamma_5)U + \frac{1}{2}(1 - \gamma_5)U^\dagger, \quad (3)$$

where  $U = \exp\{i(\tau_i \pi_i + \lambda_a K_a + \lambda_8 \eta + \eta_0)\}$ .

In Ref.[11] this Lagrangian has been applied to study  $\tau$  mesonic decays systematically. Theoretical results agree well with data. In this paper the Lagrangian(2) is used to predict all the 10 coefficients of ChPT.

As a matter of fact, in Ref.[9] two of the coefficients have been determined from the contribution of the  $\rho$  resonance in  $\pi\pi$  scattering. There are contact terms which make less contribution. In this paper a complete expressions of the coefficients  $L_{1,2,3}$  are presented.

The vertex  $\rho\pi\pi$  is[9]

$$\begin{aligned}\mathcal{L}^{\rho\pi\pi} &= f_{\rho\pi\pi}(q^2)\epsilon_{ijk}\rho_\mu^i\pi_j\partial^\mu\pi_k, \\ f_{\rho\pi\pi}(q^2) &= \frac{2}{g}\{1 + \frac{q^2}{2\pi^2 f_\pi^2}[(1 - \frac{2c}{g})^2 - 4\pi^2 c^2]\}, \\ c &= \frac{f_\pi^2}{2gm_\rho^2},\end{aligned}\tag{4}$$

where  $g$  is the universal coupling constant in this effective theory[9] and  $q$  is the momentum of the  $\rho$  meson. The effective Lagrangian of  $\pi\pi$  scattering at low energy ( $q^2 < m_\rho^2$ ) is derived from Eq.(4) and Eq.(13) of Ref.[9]

$$\begin{aligned}\mathcal{L} &= \frac{1}{2f_\pi^2}\partial_\mu\pi^2\partial^\mu\pi^2 \\ &+ \frac{16}{f_\pi^4}\{[\frac{1}{2}\frac{1}{(4\pi)^2}(1 - \frac{2c}{g})^2(-1 + \frac{4c}{g} + \frac{12c^2}{g^2}) - \frac{c^4}{g^2}]\partial_\mu\pi_i\partial^\mu\pi_i\partial_\nu\pi_j\partial^\nu\pi_j \\ &+ [\frac{1}{(4\pi)^2}(1 - \frac{2c}{g})^2(1 - \frac{4c}{g} - \frac{4c^2}{g^2}) + \frac{c^4}{g^2}]\partial_\mu\pi_i\partial^\nu\pi_i\partial_\mu\pi_j\partial^\nu\pi_j\} \\ &- \frac{4}{f_\pi^4}(1 - \frac{2c}{g})\frac{2c}{g}\{2gc + \frac{1}{\pi^2}(1 - \frac{2c}{g})\}\{\partial_\mu\pi_i\partial^\mu\pi_i\partial_\nu\pi_j\partial^\nu\pi_j - \partial_\mu\pi_i\partial^\nu\pi_i\partial_\mu\pi_j\partial^\nu\pi_j\}.\end{aligned}\tag{5}$$

Comparing with Lagrangian of the ChPT(1), we obtain

$$\begin{aligned}
2(L_1 + L_2) + L_3 &= \frac{1}{4} \frac{1}{(4\pi)^2} \left(1 - \frac{2c}{g}\right)^4 \\
L_2 &= \frac{1}{4} \frac{c^4}{g^2} + \frac{1}{4} \frac{1}{(4\pi)^2} \left(1 - \frac{2c}{g}\right)^2 \left(1 - \frac{4c}{g} - \frac{4c^2}{g^2}\right) \\
&+ \frac{1}{8} \left(1 - \frac{2c}{g}\right) \frac{c}{g} \left\{2gc + \frac{1}{\pi^2} \left(1 - \frac{2c}{g}\right)\right\}.
\end{aligned} \tag{6}$$

Only two of the three coefficients are determined from  $\pi\pi$  scattering. Therefore, another process is needed. We choose  $\pi K$  scattering[12] at low energy to do the job.  $\rho$  and  $K^*$  resonances contribute to  $\pi K$  scattering. Using Eq.(13) in Ref.[9], the  $\rho\pi\pi$  vertex and

$$\mathcal{L}^{K^*\pi\pi} = f_{\rho\pi\pi}(q^2) f_{abk} K_\mu^{a*} (K_b \partial^\mu \pi_k - \pi_k \partial^\mu K_b) [11], \tag{7}$$

in the chiral limit the two isospin amplitudes of  $\pi K$  scattering at low energy are derived as

$$\begin{aligned}
T^{\frac{3}{2}} &= -\frac{2}{f_\pi^2} s \\
&+ \frac{16}{f_\pi^4} \left\{ \frac{1}{(4\pi)^2} \frac{4c^2}{g^2} \left(1 - \frac{2c}{g}\right)^2 - \frac{c^4}{2g^2} \right\} \{t^2 + u^2\} \\
&+ \frac{16}{f_\pi^4} \left\{ \frac{1}{(4\pi)^2} \left(1 - \frac{2c}{g}\right)^2 \left(1 - \frac{4c}{g} - \frac{4c^2}{g^2}\right) + \frac{c^4}{g^2} \right\} s^2 \\
&- \frac{1}{f_\pi^4} \frac{4c}{g} \left(1 - \frac{2c}{g}\right) \left\{2gc + \frac{1}{\pi} \left(1 - \frac{2c}{g}\right)\right\} \{u(s-t) + t(s-u)\}, \\
T^{\frac{1}{2}} &= -\frac{1}{f_\pi^2} (3u - s) \\
&+ \frac{16}{f_\pi^4} \left\{ \frac{1}{(4\pi)^2} \frac{4c^2}{g^2} \left(1 - \frac{2c}{g}\right)^2 - \frac{c^4}{2g^2} \right\} t^2 \\
&- \frac{8}{f_\pi^4} \left\{ \frac{1}{(4\pi)^2} \left(1 - \frac{2c}{g}\right)^2 \left(1 - \frac{4c}{g} - \frac{16c^2}{g^2}\right) + \frac{5}{2} \frac{c^4}{g^2} \right\} s^2
\end{aligned}$$

$$\begin{aligned}
& + \frac{8}{f_\pi^4} \left\{ \frac{1}{(4\pi)^2} \left(1 - \frac{2c}{g}\right)^2 \left(3 - \frac{12c}{g} - \frac{16c^2}{g^2}\right) + \frac{7}{2} \frac{c^4}{g^2} \right\} u^2 \\
& + \frac{1}{f_\pi^4} \frac{2c}{g} \left(1 - \frac{2c}{g}\right) \left\{ 2gc + \frac{1}{\pi} \left(1 - \frac{2c}{g}\right) \right\} \{-3s(u-t) + u(s-t) + 4t(s-u)\}, \quad (8)
\end{aligned}$$

where

$$s = (p_1 + p_2)^2, \quad t = (p_1 - p_3)^2, \quad u = (p_1 - p_4)^2,$$

$p_{1,3}$  are the momenta of initial and final pions and  $p_{2,4}$  are the momenta of initial and final kaons respectively. The two isospin amplitudes of  $\pi K$  scattering can be derived from the chiral perturbation theory(1) too. Comparing the partial waves obtained from Eqs.(8) and ChPT, besides Eqs.(6) obtained from  $\pi\pi$  scattering we obtain

$$\begin{aligned}
-2L_1 + L_2 &= 0, \\
L_3 &= -\frac{3}{16} \frac{2c}{g} \left(1 - \frac{2c}{g}\right) \left\{ 2gc + \frac{1}{\pi^2} \left(1 - \frac{2c}{g}\right) \right\} - \frac{1}{2} \frac{1}{(4\pi)^2} \left(1 - \frac{2c}{g}\right)^2 \left(1 - \frac{4c}{g} - \frac{8c^2}{g^2}\right) - \frac{3}{4} \frac{c^4}{g^2}, \\
L_1 &= \frac{1}{32} \frac{2c}{g} \left(1 - \frac{2c}{g}\right) \left\{ 2gc + \frac{1}{\pi^2} \left(1 - \frac{2c}{g}\right) \right\} + \frac{1}{8} \frac{1}{(4\pi)^2} \left(1 - \frac{2c}{g}\right)^2 \left(1 - \frac{4c}{g} - \frac{4c^2}{g^2}\right) + \frac{1}{4} \frac{c^4}{g^2}. \quad (9)
\end{aligned}$$

The first equation of Eq.(9) has been obtained in Ref.[8].

It is learned from Ref.[9] that  $g^2 \sim O(N_C)$ ,  $f_\pi^2 \sim O(N_C)$ , and  $m_\rho^2 \sim O(1)$  in large  $N_C$  expansion. Therefore,  $L_{1,2,3} \sim O(N_C)$ . The predictions of  $L_{1-3}$  are made at the tree level, therefore, the Eq.(9) actually means that  $-2L_1 + L_2 \sim O(1)$  in large  $N_C$  expansion.

According to Ref.1(b), the coefficients  $L_{4-8}$  are determined from the quark mass expansions of  $m_\pi^2$ ,  $m_K^2$ ,  $m_\eta^2$ ,  $f_\pi$ ,  $f_K$ , and  $f_\eta$ . Therefore, the quark mass term of the Lagrangian(5)

is needed to be taken into account in deriving the effective Lagrangian of mesons. The expressions of the masses and the decay constants of the pseudoscalars are found from the real part of the effective Lagrangian. Using the method presented in Ref.[9], in Euclidean space the real part of the effective Lagrangian of mesons with quark masses is written as

$$\begin{aligned}\mathcal{L}_{RE} = & \frac{1}{2} \int d^D x \frac{d^D p}{(2\pi)^D} \sum_{n=1}^{\infty} \frac{1}{n} \frac{1}{(p^2 + m^2)^n} \\ & Tr\{(\gamma \cdot \partial - i\gamma \cdot v + i\gamma \cdot a\gamma_5)(\gamma \cdot \partial - i\gamma \cdot v - i\gamma \cdot a\gamma_5) + 2ip \cdot (\partial - iv - ia) \\ & + m\gamma \cdot Du - i[\gamma \cdot v, M] + i\{\gamma \cdot a, M\}\gamma_5 - m(\hat{u}M + Mu) - M^2\}^n.\end{aligned}\quad (10)$$

where  $D_\mu u = \partial_\mu u - i[v_\mu, u] + i\{a_\mu, u\}\gamma_5$  and  $\hat{u} = \exp\{-i\gamma_5[\tau_i\pi_i + \lambda_a K_a + \lambda_8\eta_8 + \eta_0]\}$ . Comparing with Eq.(11) of Ref.[9], there are new terms in which the quark mass matrix  $M$  is involved. As done in Ref.[9], the mesons fields in Eq.(10) have to be normalized.

The masses of the pseudoscalar mesons are derived from Eq.(10). The masses of the pseudoscalars can be calculated to any order in quark masses. Up to the second order in quark masses we obtain

$$\begin{aligned}m_{\pi^\pm}^2 &= \frac{4}{f_{\pi^\pm}^2} \left\{ -\frac{1}{3} \langle \bar{\psi}\psi \rangle (m_u + m_d) - \frac{F^2}{4} (m_u + m_d)^2 \right\}, \\ m_{\pi^0}^2 &= \frac{4}{f_{\pi^0}^2} \left\{ -\frac{1}{3} \langle \bar{\psi}\psi \rangle (m_u + m_d) - \frac{F^2}{2} (m_u^2 + m_d^2) \right\}, \\ m_{K^+}^2 &= \frac{4}{f_{K^+}^2} \left\{ -\frac{1}{3} \langle \bar{\psi}\psi \rangle (m_u + m_s) - \frac{F^2}{4} (m_u + m_s)^2 \right\}, \\ m_{K^0}^2 &= \frac{4}{f_{K^0}^2} \left\{ -\frac{1}{3} \langle \bar{\psi}\psi \rangle (m_d + m_s) - \frac{F^2}{4} (m_d + m_s)^2 \right\},\end{aligned}$$

$$m_{\eta_8}^2 = \frac{4}{f_{\eta_8}^2} \left\{ -\frac{1}{3} \langle \bar{\psi}\psi \rangle + \frac{1}{3} (m_u + m_d + 4m_s) - \frac{F^2}{6} (m_u^2 + m_d^2 + 4m_s^2) \right\}, \quad (11)$$

where  $F^2$  is defined in Ref.[9] as

$$F^2 = \frac{N_C}{\pi^4} m^2 \int \frac{d^4 p}{(p^2 + m^2)^2},$$

$\langle \bar{\psi}\psi \rangle$  is the quark condensate of three flavors. In terms of a cut-off  $\Lambda$  the quark condensate is expressed as (using Eqs.(40) and (43) of Ref.[9])

$$\langle \bar{\psi}\psi \rangle = \frac{i}{(2\pi)^4} \text{Tr} \int d^4 p \frac{\gamma \cdot p - m}{p^2 - m^2} = \frac{m^3 N_C}{4\pi^2} \left\{ \frac{\Lambda^2}{m^2} - \log\left(\frac{\Lambda^2}{m^2} + 1\right) \right\}. \quad (12)$$

The universal coupling constant  $g$  is expressed as [9]

$$g^2 = \frac{F^2}{6m^2} = \frac{1}{2\pi^2} \left\{ \log\left(\frac{\Lambda^2}{m^2} + 1\right) + \frac{1}{\frac{\Lambda^2}{m^2} + 1} - 1 \right\}. \quad (13)$$

The decay constants of the pseudoscalars,  $f_\pi$ ,  $f_K$ , and  $f_\eta$ , are defined by normalizing the pseudoscalar fields as done in Refs.[9,10]. In this paper we calculate the contributions of the quark masses to these decay constants to  $O(m_q)$ . Using the effective Lagrangian(10), the decay constants can be calculated to any order in quark masses. As indicated in Refs.[9,10], there is mixing between the axial-vector field and corresponding pseudoscalar field. The mixing results in the shifting of the axial-vector field  $a_\mu$

$$a_\mu \rightarrow \frac{1}{g_a} a_\mu - \frac{c_a}{g_a} \partial_\mu P, \quad (14)$$



where  $P$  is the corresponding pseudoscalar field,  $g_a$  is the normalization constant of the  $a_\mu$  field, and  $c_a$  is the mixing coefficient. Both  $g_a$  and  $c_a$  are determined in the chiral limit in Ref.[9]. For pion and  $a_1$  fields it is obtained by eliminating the mixing between pion and  $a_1$  fields

$$\frac{c_a}{g_a} = \frac{1}{g_a^2 m_a^2} \left\{ \frac{F^2}{2} + \left( \frac{F^2}{8m^2} - \frac{3}{4\pi^2} \right) 2m(m_u + m_d) \right\}, \quad (15)$$

where  $m_a$  is the mass of the  $a_1$  meson, which is determined from the Lagrangian(10) as

$$g_a^2 m_a^2 = F^2 + g^2 m_\rho^2 + 6g_{0a}^2 m(m_u + m_d), \quad (16)$$

where  $g_{0a}^2$  is expressed as[9]

$$g_{0a}^2 = g^2 \left( 1 - \frac{1}{2\pi^2 g^2} \right). \quad (17)$$

Up to the first order in quark masses, the decay constant  $f_\pi^2$  is obtained from the Lagrangian(10)

$$f_\pi^2 = f_{\pi 0}^2 \left\{ 1 + f \frac{m_u + m_d}{m} \right\}, \quad (18)$$

where

$$f_{\pi 0}^2 = F^2 \left( 1 - \frac{2c}{g} \right) [9] \quad (19)$$

and

$$f = \left( 1 - \frac{2c}{g} \right) \left( 1 - \frac{1}{2\pi^2 g^2} \right) - 1 + \frac{4}{\pi^2 f_{\pi 0}^4} \left( -\frac{1}{3} \right) < \bar{\psi}\psi > m \left( 1 - \frac{2c}{g} \right) \left( 1 - \frac{c}{g} \right). \quad (20)$$

In the same way,  $f_{K^+}^2$ ,  $f_{K^0}^2$ , and  $f_{\eta_8}^2$  are found

$$f_{K^+}^2 = f_{\pi^0}^2 \left\{ 1 + f \frac{m_u + m_s}{m} \right\}, \quad (21)$$

$$f_{K^0}^2 = f_{\pi^0}^2 \left\{ 1 + f \frac{m_d + m_s}{m} \right\}, \quad (22)$$

$$f_{\eta_8}^2 = f_{\pi^0}^2 \left\{ 1 + \frac{1}{3} f \frac{m_u + m_d + 4m_s}{m} \right\}. \quad (23)$$

Substituting Eqs.(18,21,22,23) into Eq.(11), to the second order in quark masses the masses of the pseudoscalar mesons are obtained

$$\begin{aligned} m_{\pi^\pm}^2 &= \frac{4}{f_{\pi^0}^2} \left\{ -\frac{1}{3} \langle \bar{\psi}\psi \rangle (m_u + m_d) - \frac{F^2}{4} (m_u + m_d)^2 + \frac{f}{3} \langle \bar{\psi}\psi \rangle \frac{1}{m} (m_u + m_d)^2 \right\}, \\ m_{\pi^0}^2 &= \frac{4}{f_{\pi^0}^2} \left\{ -\frac{1}{3} \langle \bar{\psi}\psi \rangle (m_u + m_d) - \frac{F^2}{2} (m_u^2 + m_d^2) + \frac{f}{3} \langle \bar{\psi}\psi \rangle \frac{1}{m} (m_u + m_d)^2 \right\}, \\ m_{K^+}^2 &= \frac{4}{f_{\pi^0}^2} \left\{ -\frac{1}{3} \langle \bar{\psi}\psi \rangle (m_u + m_s) - \frac{F^2}{4} (m_u + m_s)^2 + \frac{f}{3} \langle \bar{\psi}\psi \rangle \frac{1}{m} (m_u + m_s)^2 \right\}, \\ m_{K^0}^2 &= \frac{4}{f_{\pi^0}^2} \left\{ -\frac{1}{3} \langle \bar{\psi}\psi \rangle (m_d + m_s) - \frac{F^2}{4} (m_d + m_s)^2 + \frac{f}{3} \langle \bar{\psi}\psi \rangle \frac{1}{m} (m_d + m_s)^2 \right\}, \\ m_{\eta_8}^2 &= \frac{4}{f_{\pi^0}^2} \left\{ -\frac{1}{3} \langle \bar{\psi}\psi \rangle \frac{1}{3} (m_u + m_d + 4m_s) - \frac{F^2}{4} \frac{2}{3} (m_u^2 + m_d^2 + 4m_s^2) \right. \\ &\quad \left. + \frac{f}{3} \langle \bar{\psi}\psi \rangle \frac{1}{9} \frac{1}{m} (m_u + m_d + 4m_s)^2 \right\} \\ &= \frac{4}{f_{\pi^0}^2} \left\{ -\frac{1}{3} \langle \bar{\psi}\psi \rangle \frac{1}{3} (m_u + m_d + 4m_s) + \left( \frac{f}{m} \frac{1}{3} \langle \bar{\psi}\psi \rangle - \frac{F^2}{4} \right) \frac{1}{9} (m_u + m_d + 4m_s)^2 \right. \\ &\quad \left. - \frac{2F^2}{9} \left[ \frac{1}{2} (m_u + m_d) - m_s \right]^2 - \frac{F^2}{12} (m_d - m_u)^2 \right\}. \end{aligned} \quad (24)$$

The mass formulas(24) are the same as the ones presented in Ref.(1(b)). The mass difference

of charged and neutral pions is found from Eqs.(24)

$$m_{\pi^\pm}^2 - m_{\pi^0}^2 = (1 - \frac{2c}{g})^{-1}(m_d - m_u)^2. \quad (25)$$

Comparing with Eqs.(24) with the Eqs.(10.7,10.8) of Ref.(1(b)), following coefficients of the ChPT are predicted

$$L_4 = 0, \quad L_6 = 0, \quad (26)$$

$$L_5 = \frac{f_{\pi^0}^2 f}{8mB_0}, \quad (27)$$

$$L_8 = -\frac{F^2}{16B_0^2}, \quad (28)$$

$$3L_7 + L_8 = -\frac{F^2}{16B_0^2}, \quad L_7 = 0, \quad (29)$$

where

$$B_0 = \frac{4}{f_{\pi^0}^2}(-\frac{1}{3}) < \bar{\psi}\psi >. \quad (30)$$

Using Eq.(30),  $L_5$  and  $L_8$  are written as

$$L_5 = \frac{1}{32Q}(1 - \frac{2c}{g})\{(1 - \frac{2c}{g})^2(1 - \frac{1}{2\pi^2 g^2}) - (1 - \frac{2c}{g}) + \frac{4}{\pi^2}Q(1 - \frac{c}{g})\}, \quad (31)$$

$$L_8 = -\frac{1}{1536g^2 Q^2}(1 - \frac{2c}{g})^2, \quad (32)$$

where

$$Q = -\frac{1}{108g^4} \frac{1}{m^3} < \bar{\psi}\psi >. \quad (33)$$

The Eqs.(12,13) show that  $Q$  is a function of the universal coupling constant  $g$  only. By fitting  $\rho \rightarrow e^+e^-$ ,  $g$  is determined to be 0.39[11]. The numerical value of  $Q$  is 4.54.

Both  $L_5$  and  $L_8$  are at  $O(N_C)$ .  $L_4$ ,  $L_6$ , and  $L_7$  are from loop diagrams of mesons and are at  $O(1)$  in large  $N_C$  expansion.

According to Refs.[1(b),3],  $L_9$  and  $L_{10}$  are determined by  $\langle r^2 \rangle_\pi$  and the amplitudes of pion radiative decay,  $\pi^- \rightarrow e^- \gamma \nu$

$$L_9 = \frac{f_\pi^2}{48} \langle r_\pi^2 \rangle, \quad (34)$$

$$L_9 = \frac{1}{32\pi^2} \frac{R}{F^V}, \quad (35)$$

$$L_{10} = \frac{1}{32\pi^2} \frac{F^A}{F^V} - L_9, \quad (36)$$

where  $R$ ,  $F^V$ , and  $F^A$  are  $r_A$ ,  $h_V$ , and  $h_A$  of Ref.[3] respectively. In Ref.[11] we have derived the expression of pion radius as

$$\langle r^2 \rangle_\pi = \frac{6}{m_\rho^2} + \frac{3}{\pi^2 f_\pi^2} \left\{ \left(1 - \frac{2c}{g}\right)^2 - 4\pi^2 c^2 \right\} \quad (37)$$

which agrees with the data very well. Using Eqs.(34,37), it is predicted

$$L_9 = \frac{1}{4}cg + \frac{1}{16\pi^2} \left\{ \left(1 - \frac{2c}{g}\right)^2 - 4\pi^2 c^2 \right\} [14], \quad (38)$$

On the other hand,  $L_9$  is determined by Eq.(35) too. The three form factors of the decay amplitude of  $\pi^- \rightarrow e^- \gamma \nu$  are presented in our paper[11]

$$F^V = \frac{m_\pi}{2\sqrt{2}\pi^2 f_\pi}, \quad (39)$$

$$F^A = \frac{1}{2\sqrt{2}\pi^2} \frac{m_\pi}{f_\pi} \frac{m_\rho^2}{m_a^2} \left(1 - \frac{2c}{g}\right) \left(1 - \frac{1}{2\pi^2 g^2}\right)^{-1}, \quad (40)$$

$$R = \frac{g^2}{\sqrt{2}} \frac{m_\pi}{f_\pi} \frac{m_\rho^2}{m_a^2} \left\{ \frac{2c}{g} + \frac{1}{\pi^2 g^2} \left(1 - \frac{2c}{g}\right) \right\} \left(1 - \frac{1}{2\pi^2 g^2}\right)^{-1} + \sqrt{2}cg \frac{m_\pi}{f_\pi} [15]. \quad (41)$$

Using the mass formula of the  $a_1$  meson(in the chiral limit)[9]

$$\left(1 - \frac{1}{2\pi^2 g^2}\right) m_a^2 = \frac{F^2}{g^2} + m_\rho^2,$$

it is obtained

$$\frac{m_\rho^2}{m_a^2} \left(1 - \frac{1}{2\pi^2 g^2}\right)^{-1} = 1 - \frac{2c}{g}. \quad (42)$$

Substituting Eq.(42) into R(41), it is derived

$$R = \frac{1}{3\sqrt{2}} m_\pi f_\pi < r^2 >_\pi. \quad (43)$$

This is the expression obtained by applying PCAC to this process[14]. Therefore, the coefficient  $L_9$  derived from Eq.(35) is the same as the one obtained from Eq.(34).

Using Eq.(42), the ratio  $\frac{F^A}{F^V}$  is written as

$$\frac{F^A}{F^V} = \left(1 - \frac{2c}{g}\right)^2. \quad (44)$$

The coefficient  $L_{10}$  is found from Eq.(36)

$$L_{10} = -\frac{1}{4}cg + \frac{1}{4}c^2 - \frac{1}{32\pi^2} \left(1 - \frac{2c}{g}\right)^2. \quad (45)$$

Both  $L_9$  and  $L_{10}$  are at  $O(N_C)$ .

In the expressions of the coefficients(6,9,26,29,31,32,38,45) there are two parameters:  $g$  and  $f_\pi^2/m_\rho^2$  which have been determined already. The coefficients of the ChPT are completely determined by this effective chiral theory. The numerical values of the 10 coefficients predicted by present theory are listed in table 3.

To summarize the results. It is necessary to point out that theoretical results of the processes,  $\pi\pi$  scattering,  $\pi K[16]$  scattering, masses of the pseudoscalar mesons, the radius of pion, and the form factors of  $\pi \rightarrow \gamma e \nu$  have been tested. The orders of the coefficients of the ChPT in large  $N_C$  expansion are predicted as  $L_{1,2,3,5,8,9,10} \sim O(N_C)$  and  $L_{4,6}$  are  $O(1)$ . These predictions agree with Ref.[1(b)]. It is also predicted that  $L_7 \sim O(1)$  in this paper. It is necessary to point out that the expressions of the coefficients  $L_{4,5,6,7,8}$  are derived from Eq.(24), however, Eq.(11) is used to fit the masses of the pseudoscalar mesons. It is found that this way leads to consistent determination of the quark masses. The results will be presented somewhere else. There are no new parameters in predicting the coefficients of the ChPT.

The author wishes to thank B.Holstein for discussion. This research was partially supported by DOE Grant No. DE-91ER75661.

## References

- [1] J.Gasser and H.Leutwyler, (a) Ann.Phys.(N.Y.)**58**,(1984)142;(b) Nucl.Phys.  
**B250**, (1985)465; (c)**B250**, (1985)517.
- [2] J.Balog, Phys.Lett. **149B**, (1984)197; P.Simic, Phys.Rev.Lett. **55**, (1985)40; D.Ebert and  
H.Reinhardt, Nucl.Phys. **B271**, (1986)188; R.E.Gamboa-Saravi et al., Ann Phys. **157**,  
(1984)369; S.-K.Hu et al., Phys.Rev. **D30**, (1984)836.
- [3] J.F.Donoghue and B.R.Holstein, Phys.Rev.**D40**, (1989)2378.
- [4] J.F.Donoghue, C.Ramirez and G.Valencia, Phys.Rev. **D39**, (1989)1947; G.Ecker,  
J.Gasser, A.Pich, and E.de Rafael, Nucl. Phys. **B223**, (1989)425.
- [5] D.Ebert and M.K.Volkov, Phys.Lett. **101B**, (1981)252; V.Bernard et al., Phys.Lett.  
**B205**, (1988)16; G.Valencia and M.Praszalowicz, Nucl.Phys. **B341**, (1990)27.
- [6] D.Espriu, E.de Rafael and J.Taron, Nucl.Phys. **B345**, (1990)22, errat. **B355**, (1991)278.
- [7] B.Holdom, J.Terning and K.Verbeek, Phys.Lett. **B232**, (1989)351.
- [8] (a)G.Ecker, J.Gasser, H.Leutwyler, A.Pich, and E.Rafael, Phys. Lett. **B223**, (1989)425;  
(b)J.Bijnens, Nucl.Phys. **B390**, (1993)501.
- [9] B.A.Li, Phys.Rev., **D52**, (1995)5165.
- [10] B.A.Li, Phys.Rev., **D52**, (1995)5184.

- [11] B.A.Li, Phys.Rev., **D55**,(1997)1425, (1997)1436.
- [12] V.Bernard, N.Kaiser and U.G.Meissner, Phys.Rev.**D43**, R2757(1991), Nucl.Phys. **B357**,(1991)129, **B364**,(1991)283; A.Dobado and J.R.Peaez, Phys.Rev.**D47**,(1993)4883; L.Beldjoud1 and T.N.Truong, Phys.Lett.**B351**,(1995)357; J.Sa Borges and F.R.A.Simao, Phys.Rev.**D53**,(1996)4806.
- [13] In Eq.(67) of Ref.[19] the factor  $\frac{1}{2}$  should be taken off and there should be an additional term from the form factor  $\frac{m_\rho^2}{m_\rho^2 - k^2}$ .
- [14] This expression is obtained by B.Holstein first, private communication.
- [15] A.Kersch, Nucl.Phys. **B263**,(1986)475.
- [16] B.A.Li, D.N.Gao, and M.L.Yan, in preparation.

Table 1: Values of the coefficients[1(b)]

| $10^3 L_1$  | $10^3 L_2$   | $10^3 L_3$     | $10^3 L_4$  | $10^3 L_5$   | $10^3 L_6$   | $10^3 L_7$    | $10^3 L_8$   | $10^3 L_9$   | $10^3 L_{10}$ |
|-------------|--------------|----------------|-------------|--------------|--------------|---------------|--------------|--------------|---------------|
| $.9 \pm .3$ | $1.7 \pm .7$ | $-4.4 \pm 2.5$ | $0. \pm .5$ | $2.2 \pm .5$ | $0. \pm 0.3$ | $-.4 \pm .15$ | $1.1 \pm .3$ | $7.4 \pm .7$ | $-6. \pm .7$  |



Table 2: Coefficients obtained by models

|                                         | Vectors[4] | Quark[2] | Ref.[6] | Ref.[7] | Nucleon loop | Linear $\sigma$ model | ENJL |
|-----------------------------------------|------------|----------|---------|---------|--------------|-----------------------|------|
| $(L_1 + \frac{1}{2}L_3) \times 10^{-3}$ | -2.1       | -.8      | 2.1     | 1.1     | -.8          | -.5                   |      |
| $L_1 \times 10^{-3}$                    |            |          |         |         |              |                       | 0.8  |
| $L_2 \times 10^{-3}$                    | 2.1        | 1.6      | 1.6     | 1.8     | .8           | 1.5                   | 1.6  |
| $L_3 \times 10^{-3}$                    |            |          |         |         |              |                       | -4.1 |
| $L_4 \times 10^{-3}$                    |            |          |         |         |              |                       | 0.   |
| $L_5 \times 10^{-3}$                    |            |          |         |         |              |                       | 1.5  |
| $L_6 \times 10^{-3}$                    |            |          |         |         |              |                       | 0.   |
| $L_7 \times 10^{-3}$                    |            |          |         |         |              |                       |      |
| $L_8 \times 10^{-3}$                    |            |          |         |         |              |                       | 0.8  |
| $L_9 \times 10^{-3}$                    | 7.3        | 6.3      | 6.7     | 6.1     | 3.3          | .9                    | 6.7  |
| $L_{10} \times 10^{-3}$                 | -5.8       | -3.2     | -5.8    | -5.2    | -1.7         | -2.0                  | -5.5 |

Table 3: Predictions of the Values of the coefficients

| $10^3 L_1$ | $10^3 L_2$ | $10^3 L_3$ | $10^3 L_4$ | $10^3 L_5$ | $10^3 L_6$ | $10^3 L_7$ | $10^3 L_8$ | $10^3 L_9$ | $10^3 L_{10}$ |
|------------|------------|------------|------------|------------|------------|------------|------------|------------|---------------|
| 1.0        | 2.0        | -5.16      | 0          | 4.77       | 0          | 0          | -0.079     | 8.3        | -7.1          |